

# DID PREVALENCE, DIAGNOSIS AND KEY INDICATORS

—by Myron Walters, MA, CMT, LMFT

Dissociative Identity Disorder (DID), previously called Multiple Personality Disorder, is typically thought of as a fairly rare disorder. Prevalence studies in multiple countries have indicated otherwise, with estimates ranging from 1 to 3.1% across the general population, and 3.3 to 14% across clinical populations (Richardson, 2009, pp. 19–23).

Doing my own reality check on this, I've seen just over 100 clients in my career thus far, with 8 fairly certain DID diagnoses and two more probable DID diagnoses. This is about 8 to 10% of my clinical population, well within the 3.3 to 14% previously reported.

DID clients typically have 6 to 7 psychiatric diagnoses before they get diagnosed with DID (Ross, 1997, p. 122)—in order of prevalence: mood disorders, personality disorders, anxiety disorders, schizophrenia, substance abuse, adjustment disorder, somatization disorder, eating disorder and an organic mental disorder. In other words, if a client, or potential client, has many diagnoses, especially with a number of failed therapies, it would be prudent to get interested in the possibility of a dissociative disorder being present.

Unfortunately, most psychiatrists tend not to diagnose DID—psychiatry still seems to be recovering from Freud's 1905 public rejection of the seduction theory, which linked dissociative symptoms to sexual abuse (Richardson, 2009, p. 5). It also doesn't help that DID is not a mental health parity diagnosis, whereas schizophrenia, bipolar and depression are parity diagnoses.

Diagnosing DID is not easy, even for the clinician experienced in dissociative disorders. A rigorous workup is required. Many times it is not possible to make a quick diagnosis because the client is intentionally or unintentionally hiding the key parameters needed for a clinician to make the diagnosis. In many respects, the functional purpose of DID is to hide from view. This makes sense from and an etiology point of view, when one considers the high degree of correlation of DID with childhood abuse (physical, sexual, emotional and neglect) (Putnam, 1989, pp. 46–50).

The gold standard for diagnosing DID is the Structured Clinical Interview for DSM-IV Dissociative Disorders, otherwise known as SCID-D (Steinberg & et. al., 1993), but this takes about 3 ½ hours for a trained clinician to administer.

I can also highly recommend Lowenstein's excellent and detailed office mental status for dissociative symptoms and DID (Lowenstein, 1991).

A rapid screening, self-report tool for a dissociative disorder is the Dissociative Experience Scale (DES) (Carlson & Putnam, 1993). Although not perfect, this 28 question tool can give a quick indication if further investigation would be helpful.

Another self-report tool that I've found useful is the Multidimensional Inventory of Dissociation v.6.0 (MID-6) (Dell, 2006). This has 218 questions and gives a dissociative mapping for the client.

The MID-6 and DES are both available from the International Society for the Study of Trauma and Dissociation (ISSTD).

The presence of two or more distinct identities or personality states is the key criterion for DID in the DSM-IV-TR (APA, 2000). Unfortunately, this key criterion is rarely seen in or disclosed by a client until a strong therapeutic alliance has been established. This

is one of the contributing factors to a firm DID diagnosis sometimes taking years.

Below are some of the key features I have found beneficial to check during my initial client intake. If I get a hit on a constellation of these, I definitely want to look closer for a dissociative disorder, e.g., I would then give the DES and/or MID-6.

1. Multiple previous diagnoses (see above)
2. Multiple failed previous therapies (see above)
3. Time loss or time gaps which are not accounted for by use of drugs, alcohol or medical reasons (e.g., epilepsy), especially large gaps of memory from childhood. This is actually one of the DSM-IV-TR criteria, so I give this one high importance.
4. Suicidality
5. Self-harm activities
6. Inner voices that are ego-dystonic to the client—generally, I won't learn about this though until a sufficient therapeutic alliance has been established.
7. PTSD symptoms, such as hyper vigilance, intrusive thoughts, flashbacks, etc.
8. An extensive childhood trauma history
9. A client who does a hypnotic eye-roll, where the client rolls their eyes up into the top of their head showing the whites of their eyes (see Figure 1)—this typically happens during a switch.



Figure 1: Hypnotic Eye Roll

Summarizing, because DID tends to be a hidden disorder that actually has a fairly high prevalence, wreaks havoc on the client and can result in many failed therapies, it is important for a clinician to begin to gain some awareness around when to suspect DID.

## BIBLIOGRAPHY

- APA. (2000). *Diagnostic and Statistical Manual of Mental Disorders—DSM-IV-TR*. American Psychiatric Association.
- Carlson, E. B., & Putnam, F. W. (1993). An update on the Dissociative Experience Scale. *Dissociation*, 6(1), 016-027. <http://hdl.handle.net/1794/1539>.
- Dell, P. F. (2006). The Multidimensional Inventory of Dissociation (MID): A comprehensive measure of pathological dissociation. *Journal of Trauma & Dissociation*, 7(2), 77-106; [http://dx.doi.org/10.1300/J229v07n02\\_06](http://dx.doi.org/10.1300/J229v07n02_06).
- Lowenstein, R. J. (1991). An office mental status examination for complex chronic dissociative symptoms and multiple personality disorder. *Psychiatric Clinics of North America*, 14(3), 567-604.
- Putnam, F. W. (1989). *Diagnosis and Treatment of Multiple Personality Disorder*. New York: The Guilford Press.
- Richardson, H. (2009). *An exploration of dissociation in Australia: measurement issues, predictors and phenomenology*. Swinburne University of Technology, Life and Social Sciences. Hawthorn, Victoria, Australia: Australasian Digital Theses collection; <http://hdl.handle.net/1959.3/78343>.

Ross, C. A. (1997). *Dissociative identity disorder: Diagnosis, clinical features, and Treatment of Multiple Personality*. New York: John Wiley & Sons, Inc.

Steinberg, M., & et. al. (1993). Clinical assessment of dissociative symptoms and disorders: The structured clinical interview for DSM-IV Dissociative Disorders (SCID-D). *Dissociation*(6:1), 3-15.



### About the Author



Myron Walters, MA CMT, Licensed Marriage and Family Therapist, has private practice offices in Mill Valley and Cotati. Myron works somatically, being both trauma and attachment informed. Myron's specialty is working with adults who have experienced severe childhood trauma, complex PTSD and dissociative disorders. Myron has training in trauma resolution (Somatic Experiencing), hypnosis (SCEH & ASCH), and in DID psychotherapy (ISSTD). Myron welcomes your calls and referrals at (415) 686-3445.




For those who are deficient in the Career and Vocational Counseling, Psychopharmacology or Intro to Clinical Assessment classes needed for the LPCC grandparenting requirement, there is an affordable alternative available at the University of Phoenix online. Regularly \$685/unit (a three unit class) they have discounted the class for granparenting candidates to \$250/unit. A big difference at \$750 instead of \$2055!

Contact Larissa Leon at [Larissa.Leon@Phoenix.edu](mailto:Larissa.Leon@Phoenix.edu) for further information or enrollment

## THErapy GROUPS (CONTINUED)

**PRACTICE-BUILDING OR GROUP-THERAPY CONSULT, Renee Owen, MFT** Choose one or both groups for: help creating a thriving practice—or—starting, sustaining, and successfully facilitating groups. Each group meets one Mon./month. 12:00–1:30p.m. San Rafael. 453-8117.

**LONG TERM THERAPY GROUP FOR PSYCHOTHERAPISTS, co-led by Geraldine Alpert, Ph.D. and Elaine Cooper Ph.D.** One opening. Psychodynamic/interpersonal orientation. Group currently has 3 men and 4 women. Meets in San Rafael on Wednesdays 9:30–11:00a.m. Geri Alpert 453-8969; [Geraldine.Alpert@gmail.com](mailto:Geraldine.Alpert@gmail.com)



**Care to be a mentor?**  
Just call CAMFT at 888-892-2638



### Community Institute for Psychotherapy Professional Development Series 2010-11

For full course information, CEUs, Dates, Times, and Cost, and to sign up to be a CIP Member for discounts and other benefits please visit us at [www.cipmarin.org](http://www.cipmarin.org) or call (415) 459-5999 x101

#### SEMINARS FOR PROFESSIONALS:

- **ancient zen pond, therapist jumps in ... plop! Shame Vulnerability & The Beginners' Mind** with A.Raja Hornstein, PhD, Arlene Berman, LCSW, Chris Fortin, MFT
- **Mental Health, A Vanishing but Critical Concept** presented by Nancy McWilliams, PhD
- **Emotion and Transformation: Changing Individuals, Couples and Communities** with Peter Dunlap, PhD and Jim Matto-Shepard, PhD

#### STUDY GROUPS & CLASSES:

- **Integrative Therapies: Integrating Mind/Body Therapies Into Practice** with Ilene Serlin, PhD
- **Widening the Royal Road** with Meredith Sabini, PhD, John Beebe, MD, Fariba Bogzaran, PhD
- **Dreams Consultation Group** with Meredith Sabini, PhD
- **Love in the Psychotherapy Relationship** with Bryan Wittine, PhD, Betsy Cohen, PhD, Jane Jordan, PsyD and Kirk J. Schneider, PhD
- **Treatment of the Traumatized Child** with Julie Robbins, LCSW

Please visit us on the web at [www.cipmarin.org](http://www.cipmarin.org) for more information about events and opportunities at CIP.



Community Institute for Psychotherapy

Helping individuals  
Healing families  
Strengthening  
our community

[www.cipmarin.org](http://www.cipmarin.org)  
(415) 459-5999  
[cipmarin@aol.com](mailto:cipmarin@aol.com)